

- SARVAS, R. 1962. Investigations on the flowering and seed crop of *Pinus silvestris*. Comm. Inst. Forest. Fenn. 53(4): 198 p.
- SCHUBERT, G. H., and W. J. RIETVELD. 1970. Bristlecone pine—its phenology, cone maturity and seed production in the San Francisco Peaks, Arizona. USDA Forest Serv. Res. Note RM-180, 7 p. Rocky Mountain Forest and Range Exp. Stn.
- UYEKI, H. 1927. The seeds of the genus *Pinus*, as an aid to the identification of species. Bull. Agric. and Forest. Coll., Suigen, Korea, No. 2: 129 p.
- WRIGHT, R. D. 1963. Some ecological studies on bristlecone pine in the White Mountains of California. Ph.D. Thesis: Univ. Calif., Los Angeles, 118 p.
- ZAVARIN, E., and K. SNAJBERK. 1973. Variability of the wood monoterpenoids from *Pinus aristata*. Biochem. Syst. 1:39–44.
- , K. SNAJBERK, and D. BAILEY. 1976. Variability in the essential oils of wood and foliage of *Pinus aristata* and *Pinus longaeva*. Biochem. Syst. 4:81–92.

A NEW GYPSOPHILOUS SPECIES OF PHACELIA (HYDROPHYLLACEAE) FROM COAHUILA, MEXICO

N. DUANE ATWOOD

Bureau of Land Management, Cedar City, Utah 84720

DONALD J. PINKAVA

Department of Botany and Microbiology
Arizona State University, Tempe 85281

Floristic studies of the Cuatro Ciénegas Basin in central Coahuila, Mexico, have revealed a new species of *Phacelia* growing on gypsum dunes and flats. This taxon is apparently restricted to the basin, adding to the list of gypsophiles only known from there: *Dyssodia gypsophila* Turner (1972a), *Gaillardia gypsophila* Turner (1972b), *Haploesthes robusta* I. M. Johnston (1941), *Machaeranthera gypsophila* Turner (1973a), and *M. restiformis* Turner (1973b).

Phacelia marshall-johnstonii Atwood and Pinkava, sp. nov. Plantae perennes 1.5–2.5 dm altae, caudicibus ligneis usque ad 1.5 cm diametro; caules 1–plures e basi erecti vel ascendentes ramificantes supra saepe viscidis et dense canescentes, pilis patulis 1–2 mm longis et pilis brevior et mollior 0.3–0.8 mm longis; folia aggregata diminuta sursum, petiolis 0.2–2.0 cm longis, laminis ovatis ad elliptica 0.8–5.5 cm longis 0.5–2.5 cm latis, margine grosse crenatis ad duple crenatis vel leviter lobatis, apicibus obtusis, basibus rotundatis ad subcordatis, utrinque dense hirsutis-viscidis; inflorescentiae terminales in axibus principalibus et ramis magnis cymarum compositarum scorpioidarum, cymae usque ad 13 cm longae in fructum; flores numerosi subsessiles; sepala elliptica ad oblanceolata usque ad 4.5 mm longa et usque ad 2 mm lata in fructum dense et grosse hirsuta-viscida; corollae caesiaae albidae ad extremum 5 mm longae infundibuliformia, tubis 4 mm longis glabris, lobis 1 mm longis hirsutis extis subtiliteris; antherae globosae ca 0.5 mm diametros exsertae-longae, filamentis purpureis ca 1 cm longis glabris 0.8 mm supra basim corolla insertis, appendicibus basalibus auriculiformibus 0.7 mm longis; styli longo-




FIG. 1. *Phacelia marshall-johnstonii*. a, habit; b, stem segment enlarged; c, corolla and androecium; d, calyx and gynoecium (at flowering); e, fruit in calyx; f, seed, dorsal view; g, seed, ventral view. Illustration based on holotype. Scale line = 5 mm.

exserti stamina equantes purpurei subti pubescenti grosse pilis 0.2–0.5 mm longis; capsulae ovoidae ad subglobosae 3.0–3.5 mm longae 2.5–3.0 mm latae apice subtiliter puberulentae; semina 4 elliptica 2.0–2.5 mm longa 1.1–1.3 mm lata fusca irridescentia leviter, pagina dorsali foveolata, pagina ventralis excavata reticulata, marginibus et crista centralie ventralis corrugatis leviter.

TYPE: MEXICO: COAHUILA: Cuatro Ciénegas Basin, along new road to gypsum dunes, 0.3 miles S of Poso de la Becerra, gypsum flats with *Atriplex*, *Nama*, and *Sporobolus*, 15 Aug 1975, *Reeves & Pinkava P13100* (Holotype: ASU; Isotypes: BRV, ENCB, GH, MEXU, NY, LL-TEX, UC, US).

Additional materials examined: MEXICO: COAHUILA: Poso de Escobeda, grassy banks, 17 Aug 1967, *Cole, Minckley & Pinkava P4086* (ASU); Julio's Canal, 4.5 miles SSW of Cuatro Ciénegas, along roadside, 15 Aug 1967, *Cole, Minckley & Pinkava P3812* (ASU); stabilized dunes W of headwaters of El Chiqueros, 15 Aug 1967, *Cole, Minckley & Pinkava P3953* (ASU); stabilized dunes S of Laguna Grande, 8 Jun 1968, *Lehto, Keil & Pinkava P5024* (ASU); gypsum dunes, 16 km S of Cuatro Ciénegas, ca. 2 km SW of Poso y Balneario La Becerra, 26° 52' N, 102° 09' W, 770 m, 22 Mar 1973, *Johnston, Wendt & Chiang 10334* (LL-TEX); 19 km SW of Cuatro Ciénegas, gypsum dunes, with *Petalonyx*, *Sporobolus*, *Dasyllirion*, 26° 52' N, 102° 09' W, 700 m, 11 Jun 1972, *Chiang, Wendt & Johnston 7649* (LL-TEX).

Phacelia marshall-johnstonii (Fig. 1) is apparently closely related to *P. pallida* I. M. Johnston of the *Phacelia crenulatae* group (Atwood, 1975), but is distinguished from that taxon by the densely canescent, harsh spreading hairs, only sparsely glandular pubescence and smaller seeds.

This species is named in honor of Professor Marshall C. Johnston, University of Texas, Austin, devoted scholar of the floras of Texas and the Chihuahuan Desert.

ACKNOWLEDGMENTS

We thank Dr. David Keil for critically reviewing the manuscript and Wendy Hodgson for the illustration. Field work was supported in part by National Science Foundation grants GB2461 and GB6477X awarded Dr. W. L. Minckley, Department of Zoology, Arizona State University.

LITERATURE CITED

- ATWOOD, N. D. 1975. A revision of the *Phacelia crenulatae* group (Hydrophyllaceae) for North America. *Great Basin Nat.* 35:127–190.
- JOHNSTON, I. M. 1941. New phanerogams from Mexico, IV. *J. Arnold Arb.* 22:110–124.
- TURNER, B. L. 1972a. A new species of *Dyssodia* (Compositae) from north central Mexico. *Madroño* 21:421–422.
- . 1972b. Two new gypsophilous species of *Gaillardia* (Compositae) from north central Mexico. *Southw. Nat.* 17:181–190.
- . 1973a. Two new gypsophilous species of *Machaeranthera* (Asteraceae-Astereae) from north-central Mexico. *Phytologia* 26:116–120.
- . 1973b. *Machaeranthera restiformis* (Asteraceae), a bizarre new gypsophile from north-central Mexico. *Amer. J. Bot.* 60:836–838.